

Citation for published version:

Sabater, S, Schmidt, D, Schmidt, H, Kuntze-Fechner, M, Zell, T, Isaac, C, Rajabi, N, Grieve, H, Blackaby, W, Lowe, J, Macgregor, S, Mahon, M, Radius, U & Whittlesey, M 2021, '[Ni(NHC)₂] as a scaffold for structurally characterized trans [H-Ni-PR₂] and trans [R₂P-Ni-PR₂] complexes', *Chemistry - A European Journal*, vol. 27, pp. 13221-13234. <https://doi.org/10.1002/chem.202101484>

DOI:

[10.1002/chem.202101484](https://doi.org/10.1002/chem.202101484)

Publication date:

2021

Document Version

Peer reviewed version

[Link to publication](#)

Publisher Rights

CC BY

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**[Ni(NHC)₂] as a Scaffold for Structurally Characterized *trans* [H-Ni-PR₂] and *trans*
[R₂P-Ni-PR₂] Complexes**

Sara Sabater,^{+[a]} David Schmidt,^{+[b]} Heidi Schmidt, nee Schneider,^[b] Maximilian W. Kuntze-
Fechner,^[b] Thomas Zell,^[b] Connie J. Isaac,^[a] Nasir A. Rajabi,^[c] Harry Grieve,^[a] William J. M.
Blackaby,^[a] John P. Lowe,^[a] Stuart A. Macgregor,^{+[c]} Mary F. Mahon,^[a] Udo Radius^{*[b]} and
Michael K. Whittlesey^{*[a]}

^[a] Dr. S. Sabater, C. J. Isaac, H. Grieve, Dr. W. J. M. Blackaby, Dr. J. P. Lowe, Dr. M. F.
Mahon, Prof. M. K. Whittlesey

Department of Chemistry, University of Bath, Claverton Down, Bath, BA2 7AY, UK

e-mail: m.k.whittlesey@bath.ac.uk

^[b] Dr. D. Schmidt, Dr. H. Schmidt, nee Schneider, Dr. M. W. Kuntze-Fechner, Dr. T. Zell,
Prof. Dr. U. Radius

Institut für Anorganische Chemie, Julius-Maximilians-Universität Würzburg, Am Hubland,
D-97074 Würzburg, Germany

e-mail: u.radius@uni-wuerzburg.de

Twitter: @Uni_WUE

^[c] Dr. N. A. Rajabi, Prof. S. A. Macgregor

Institute of Chemical Sciences, Heriot-Watt University, Edinburgh, EH14 4AS, UK

e-mail: S.A.Macgregor@hw.ac.uk

Twitter: @MacgregorHWU

^[+] These authors contributed equally to this work.


Supporting information (Experimental and X-ray crystallographic details, NMR spectra and
computational information) and ORCID(s) from the author(s) for this article are available on
the WWW under

Abstract

Addition of PPh_2H , PPhMeH , PPhH_2 , $\text{P}(\textit{para}\text{-Tol})\text{H}_2$, PMesH_2 and PH_3 to the two-coordinate $\text{Ni}(0)$ N-heterocyclic carbene species $[\text{Ni}(\text{NHC})_2]$ ($\text{NHC} = \text{I}^i\text{Pr}_2$, IMe_4 , IEt_2Me_2) affords a series of mononuclear, terminal phosphido Ni complexes. Structural characterisation of nine of these compounds show they exhibit unusual *trans* $[\text{H-Ni-PR}_2]$ or novel *trans* $[\text{R}_2\text{P-Ni-PR}_2]$ geometries. The bis-phosphido complexes are more accessible when smaller NHCs ($\text{IMe}_4 > \text{IEt}_2\text{Me}_2 > \text{I}^i\text{Pr}$) and phosphines are employed. P-P activation of the diphosphines $\text{R}_2\text{P-PR}_2$ ($\text{R}_2 = \text{Ph}_2, \text{PhMe}$) provides an alternative route to some of the $[\text{Ni}(\text{NHC})_2(\text{PR}_2)_2]$ complexes. DFT calculations capture these trends with P-H bond activation proceeding from unconventional phosphine adducts in which the H substituent bridges the Ni-P bond. P-P bond activation from $[\text{Ni}(\text{NHC})_2(\text{Ph}_2\text{P-PPh}_2)]$ adducts proceeds with computed barriers below 10 kcal/mol. The ability of the $[\text{Ni}(\text{NHC})_2]$ moiety to afford isolable terminal phosphido products reflects the stability of the Ni-NHC bond that prevents ligand dissociation and onward reaction.

Introduction

Isolation of the 14-electron nickel(0) bis-N-heterocyclic carbene (NHC) complex, $[\text{Ni}(\text{IMes})_2]$ (IMes = 1,3-bis(2,4,6-trimethylphenyl)imidazolin-2-ylidene), by Arduengo and co-workers in 1994 provided an early illustration of the ability of NHCs to stabilize coordinatively unsaturated transition metal complexes.^[1] In the case of $[\text{Ni}(\text{IMes})_2]$ and analogues containing even more bulky N-aryl substituents,^[2,3] however, a price is paid for this stability in the form of somewhat altered and limited reactivity.^[4,5] $[\text{Ni}(\text{NHC})_2]$ systems with higher reactivity towards small molecules^[6] are accessible through the use of N-alkyl substituted carbenes, although with the exception of $[\text{Ni}(\text{I}^t\text{Bu}_2)_2]$ (I^tBu_2 = 1,3-di-*tert*-butylimidazolin-2-ylidene)^[2b,7] these can only be generated in-situ. Particular success has been found with $[\text{Ni}_2(\text{I}^i\text{Pr}_2)_4(\text{COD})]$ (I^iPr_2 = 1,3-di-*isopropyl*imidazolin-2-ylidene)^[8] (and related molecules)^[9] as a labile source of the N-isopropyl substituted carbene complex $[\text{Ni}(\text{I}^i\text{Pr})_2]$ (**A**, Scheme 1), which is able to activate a wide range of C-X (X = F, Cl, Br),^[10,11] S-X (X = C, S)^[12] and E-H (E = Si, S) bonds.^[12,13] In a number of cases, these processes have been made catalytic, allowing the Suzuki-Miyaura cross-coupling of aryl fluorides, chlorides and bromides,^[11,14] hydrodefluorination of aryl fluorides^[15] and silane dehydropolymerisation.^[16] An even more impressive transformation builds upon the ability of **A** to cleave strained C-C bonds,^[17] allowing the coupling of 2,2'-biphenylene with $\text{PhC}\equiv\text{CPh}$ to yield 9,10-diphenylphenanthrene.^[8]



Scheme 1

Very recently, one of us reported isolation of the Ni(II) terminal phosphido phenyl complex, *trans*- $[\text{Ni}(\text{IMe}_4)_2(\text{PPh}_2)\text{Ph}]$, (**1**, Scheme 2) from the reaction of the N-methyl


substituted NHC IMe₄ (IMe₄ = 1,3,4,5-tetramethylimidazolin-2-ylidene) with the 16-electron Ni(0) complex [Ni(6-Mes)(PPh₃)₂] (6-Mes = 1,3-bis(2,4,6-trimethylphenyl)-3,4,5,6-tetrahydropyrimidin-2-ylidene). The formation of **1**, most likely through P-C activation of PPh₃ by [Ni(IMe₄)₂] (**B**),^[18] is surprising given that P-C cleavage reactions of aryl phosphines typically yield multi-metallic products with bridging phosphido ligands, rather than mononuclear complexes with terminal phosphido groups.^[19] Moreover, the isolation and structural characterization of **1** contrasts with the instability of the bis-triethylphosphine analogue, *trans*-[Ni(PEt₃)₂(PPh₂)Ph], which has been reported as being too labile even to allow spectroscopic characterization.^[20]

Our mutual interests in the bond activation reactions of **A** and **B** have now led us to conduct a joint investigation into their reactivity, together with that of the N-Et substituted derivative [Ni(IET₂Me₂)₂] (**C**; IET₂Me₂ = 1,3-diethyl-4,5-dimethylimidazolin-2-ylidene), towards P-H bonds of secondary and primary phosphines. At the outset of this work, we were aware of only a few examples of mononuclear nickel complexes with terminal phosphido ligands,^[21,22] and a single case of an isolated Ni phosphido hydrido complex, [Ni(d'bpe){P(Dmp)H}H] (d'bpe = ^tBu₂PCH₂CH₂P^tBu₂; Dmp = 2,6-Mes₂C₆H₃).^[23] Herein, we describe the use of **A-C** to generate terminal phosphido hydrido and terminal bis-phosphido complexes, nine of which have been structurally characterized.^[24] Among these are the first examples of terminal bis-phosphido complexes with *trans* R₂P-M-PR₂ geometries reported for any transition metal. DFT calculations have also been used to rationalize the formation of the terminal phosphido hydrido and terminal bis-phosphido complexes as a function of the NHCs involved.

Results and Discussion

In-situ generation of [Ni(IMe₄)₂] (B**) and [Ni(IET₂Me₂)₂] (**C**).** Given the atom inefficiency associated with the formation of **1** from [Ni(6-Mes)(PPh₃)₂] and IMe₄ (i.e. loss of both 6-Mes

and PPh₃ during reaction), a more economical pathway was afforded by simply mixing [Ni(COD)₂] with IMe₄ and PPh₃ in a 1:2.1 molar ratio at room temperature. This gave **1** (Scheme 2a) in ca. 90% yield based on NMR spectroscopy. The N-ethyl carbene derivative, *trans*-[Ni(IEt₂Me₂)₂(PPh₂)Ph] (**2**), could be prepared similarly (Scheme 2b), again presumably through activation of a P-Ph group by [Ni(IEt₂Me₂)₂] formed in-situ. The combination of the N-*i*-Pr substituted NHC I^{*i*}Pr₂Me₂ (I^{*i*}Pr₂Me₂ = 1,3-di-*isopropyl*-4,5-dimethylimidazol-2-ylidene) with [Ni(COD)₂] and PPh₃ gave a mixture of uncharacterized products. The formation of **1** and **2** was quite slow, taking days to reach completion. In the case of **2**, raising the temperature to 50 °C reduced the reaction time to just 3 h. Complex **2**, which was isolated in 45% yield, was readily characterizable by comparison of spectroscopic data to those of **1**, with a high frequency ¹³C{¹H} NMR resonance at δ = 166 ppm (d, ²*J*(C,P) = 40 Hz) diagnostic of the *ipso*-C of the diphenylphosphido ligand.



Scheme 2

The molecular structure of **2** (Figure 1) confirmed the presence of a pyramidal phosphido ligand ($\Sigma(\text{angles at P}) = 325.4^\circ$; cf. 328.5° for a tetrahedral P) and a *trans* Ph-Ni-PPh₂ geometry. In comparison to **1**, there was a slight elongation of both the Ni-C_{phenyl} and Ni-PPh₂ distances (**2**: 1.960(3) Å, 2.2851(8) Å; **1**: 1.9371(15), 2.2520(5) Å). We attribute this to

increased crowding at the nickel centre resulting from the less coplanar arrangement of the two NHC ligands in **2** (N-C-C-N dihedral angles: **2**: 142.3°, 161.6°; **1**: 159.2°, 156.7°). As will become obvious over the course of the manuscript, sterics plays a key role not only in defining the products formed between $[\text{Ni}(\text{NHC})_2]$ and specific phosphines, but also their geometries.





Figure 1. Molecular structure of *trans*- $[\text{Ni}(\text{IEt}_2\text{Me}_2)(\text{PPh}_2)\text{Ph}]$ (**2**). Ellipsoids are shown at 30% probability with all hydrogen atoms removed for clarity.

$[\text{Ni}(\text{NHC})_2(\text{PR}_2)\text{H}]$ formation with secondary phosphines. Upon addition of the secondary aryl phosphine PPh_2H to $[\text{Ni}(\text{I}^i\text{Pr})_2]$ (**A**) or $[\text{Ni}(\text{IEt}_2\text{Me}_2)_2]$ (**C**), rapid, room temperature P-H activation ensued to afford the phosphido hydrido complexes *trans*- $[\text{Ni}(\text{I}^i\text{Pr})_2(\text{PPh}_2)\text{H}]$ (**3**) and *trans*- $[\text{Ni}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_2)\text{H}]$ (**4**), respectively (Scheme 3). The *trans*- $\text{R}_2\text{P-M-H}$ geometry is rare for any transition metal,^[25] and contrasts with the *cis*- $\text{R}_2\text{P-Ni-H}$ arrangement in $[\text{Ni}(\text{d}^t\text{bpe})\{\text{P}(\text{Dmp})\text{H}\}\text{H}]$ resulting from the presence of the chelating phosphine ligand.^[23]



Scheme 3

The *trans* $\text{Ph}_2\text{P-Ni-H}$ arrangement in both **3** and **4** was established by X-ray crystallography (Figure 2) and multinuclear NMR spectroscopy. The change of ligand *trans* to phosphido group from Ph (**2**) to H (**4**) resulted in a slight shortening of the Ni- PPh_2 distance, despite the higher *trans* influence of the latter (**2**: 2.2851(8) Å; **4**: 2.2783(4) Å). This may reflect the narrowing of the $\text{C}_{\text{NHC}}\text{-Ni-C}_{\text{NHC}}$ bond angle (**2**: 175.26(12)°; **4**: 169.16(5)°) that permits a closer approach of the phosphido ligand on steric grounds. Ni-H resonances for **3** and **4** were found at δ -11.3 and δ -11.7, respectively, both with a doublet $^2J(\text{H,P})$ splitting of 73 Hz. Identification as a *trans*-coupling by comparing it with the *cis*-splitting in $[\text{Ni}(\text{d}^t\text{bpe})\{\text{P}(\text{Dmp})\text{H}\}\text{H}]^{[23a]}$ was prevented by the absence of any resolvable couplings on the Ni-H signal of the latter (even at low temperature). Although the *cis* phosphido-Ni-H splitting was likewise unresolved in the related complex $[\text{Ni}(\text{d}^t\text{bpe})\{\text{PMes}^*\text{H}\}\text{H}]$ ($\text{Mes}^* = 2,4,6\text{-}^t\text{Bu}_3\text{C}_6\text{H}_3$),^[23b] this did show a coupling of comparable large magnitude (104 Hz) coupling for $J(\text{trans phosphine-Ni-H})$.^[26] The best comparison involves the iridium bis- PH_2 hydrido complex, $[\text{Ir}(\text{PEt}_3)_2(\text{PH}_2)_2(\text{CO})\text{H}]$,^[25c] which shows large *trans* (42 Hz) and much smaller *cis* (7 Hz) values of $^2J(\text{PH}_2\text{-Ir-H})$. Final confirmation of geometry in **3** and **4** was provided by the appearance of a small (11 Hz) *cis* NHC-Ni-PPh_2 $^2J(\text{C,P})$ doublet splitting on each of the high frequency ^{13}C carbene resonances of **3** and **4**.






Figure 2. Molecular structures of (left) *trans*-[Ni(I'Pr)₂(PPh₂)H] (**3**) and (right) *trans*-[Ni(IEt₂Me₂)₂(PPh₂)H] (**4**). Ellipsoids are shown at 30% probability with all hydrogen atoms, except Ni-H, removed for clarity.

Formation of *trans*-[Ni(NHC)₂(PAr₂)₂] complexes. In contrast to the clean formation of phosphido hydrido complexes **3** and **4**,^[27] [Ni(IME₄)₂] (**B**) reacted with PPh₂H in time of mixing to give a mixture of both the phosphido hydrido complex *trans*-[Ni(IME₄)₂(PPh₂)H] (**5**) and the bis-phosphido species, *trans*-[Ni(IME₄)₂(PPh₂)₂] (**6**, Scheme 4). Conversion through to just **6** took place over ca. 24 h at room temperature.^[28]




Scheme 4

Despite this difference in initial products from **A** and **B** (Schemes 3 and 4), we found that addition of PPh_2H to $[\text{Ni}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_2)\text{H}]$ (**4**) did bring about conversion through to the bis-phosphido complex, $\text{trans}-[\text{Ni}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_2)_2]$ (**7**, Scheme 5),^[27] although the process was slow, requiring over ca.1 month at room temperature to reach completion.^[29] Steric factors likely contribute significantly to the difficulty of this transformation; indeed, exposure of the bulkier I^iPr_2 precursor **A** to even a large excess of PPh_2H failed to generate any of the bis- PPh_2 complex, $\text{trans}-[\text{Ni}(\text{I}^i\text{Pr}_2)_2(\text{PPh}_2)_2]$ (**8**), although the less bulky methylphenylphosphido derivative, $\text{trans}-[\text{Ni}(\text{I}^i\text{Pr}_2)_2(\text{PPhMe})_2]$ (**9**), could be made this way (Scheme 6).



Scheme 5

Unexpectedly, **8** could be formed through P-P bond activation of $\text{Ph}_2\text{P}-\text{PPh}_2$ in the presence of **A** (Scheme 6), which also reacted with $\text{PhMeP}-\text{PMePh}$ to give an alternative route to **9**. In fact, P-P activation proved to be a highly efficient route to bis-phosphido complexes - formation of **7** was complete within just 30 min when $\text{Ph}_2\text{P}-\text{PPh}_2$ was added to **C** and within a few hours upon addition to **4** (Scheme 5). Although P-P activation of P_4 and other cyclic/oligomeric P_x species is well-known,^[30] activation (or even simple coordination) of diphosphines at a transition metal (or *p*-block) fragment is extremely rare.^[31-33] Indeed, $\text{Ph}_2\text{P}-\text{PPh}_2$ is most commonly encountered as an unreactive elimination product in catalytic dehydrocoupling reactions of secondary phosphines.^[34]



Scheme 6

Figure 3 shows the X-ray crystal structures of **6** and **7**, which we believe represent the first structurally characterized examples of mononuclear metal complexes with *trans*-terminal phosphido ligands.^[35,36] The Ni atoms are coincident with crystallographic inversion centres in both structures, resulting in a *trans* disposition of the two sets of phenyl substituents about the P-Ni-P axis in **6** and the two sets of N-Et groups about the C-Ni-C axis in **7**. The Ni-P distance in **7** (2.2668(3) Å; 2.2628(5) Å in **6**) was shorter than in both [Ni(IEt₂Me₂)₂(PPh₂)Ph] (**2**) and [Ni(IEt₂Me₂)₂(PPh₂)H] (**4**), and there was a minor increase in the sum of the angles at the pyramidal phosphorus atoms (**2**, 325.4°; **4**, 324.3°; **7**, 329.6°).

The isolation of the NHC containing bis-phosphido complexes **6** and **7** is in marked contrast to the stabilities of the bis(triethylphosphine) derivatives, *trans*-[M(PEt₃)₂(P(SiMe₃)₂)₂].^[37] The M = Pt derivative survives long enough to allow a *trans*-R₂P-Pt-PR₂ arrangement to be confirmed spectroscopically, whereas the M = Ni analogue proved to be unstable above -70 °C. The difference in stability imparted by NHCs underlines the comparison made earlier between **1** and [Ni(PEt₃)₂(PPh₂)Ph].^[20]





Figure 3. Molecular structures of *trans*-[Ni(IME₄)₂(PPh₂)₂] (**6**) and *trans*-[Ni(IET₂Me₂)(PPh₂)₂] (**7**). Ellipsoids are shown at 30% probability and all hydrogen atoms are removed for clarity (**6**; ⁱ symmetry operation 1 – *x*, – *y*, 1 – *z* symmetry operation: **7**; ⁱ symmetry operation 1 – *x*, 1 – *y*, 1 – *z*).


Unequivocal spectroscopic evidence for the presence of *trans* R₂P-Ni-PR₂ arrangements in the four bis-phosphido complexes **6-9** was provided by (i) the small (²*J*(C,P) = 15-17 Hz) triplet splitting of the ¹³C carbenic carbon resonance and (ii) the appearance in all the spectra of a virtually coupled signal at ca. δ = 150 ppm for the magnetically inequivalent *ipso*-C of the two *trans*-phosphido ligands.

Reactivity of [Ni(NHC)₂] with primary aryl phosphines. P-H bond activation of the primary aryl phosphine PPhH₂ by **B** and **C** (Scheme 7) was also facile, although both reactions gave a mixture of the primary phosphido hydrido complexes *trans*-[Ni(NHC)₂(PPhH)H] (NHC = IMe₄ **10**, IET₂Me₂ **12**) and bis-phosphido products, *trans*-[Ni(NHC)₂(PPhH)₂] (NHC = IMe₄ **11**, IET₂Me₂ **13**). As shown in Scheme 8, **A** displayed even more complex reactivity with 1-4 equiv PPhH₂ (as well as P(*para*-Tol)H₂) giving a mixture of

three products, even when the reactions were conducted at $-30\text{ }^{\circ}\text{C}$.^[38] The formation of *trans*- $[\text{Ni}(\text{I}^i\text{Pr}_2)_2(\text{PArH})\text{H}]$ (Ar = Ph **14**, *para*-Tol **15**) was signified by the appearance of low frequency signals (vide infra) in both the ^1H and ^{31}P NMR spectra, although these complexes defied all efforts at isolation. Repeating the reactions in the presence of an alkene as a hydrogen acceptor allowed isolation and characterisation of the second component of the reactions, namely the bis-phosphido complexes, *trans*- $[\text{Ni}(\text{I}^i\text{Pr}_2)_2(\text{PArH})_2]$ (Ar = Ph **16**, *para*-Tol **17**). The diphosphene complexes, $[\text{Ni}(\text{I}^i\text{Pr}_2)_2(\eta^2\text{-ArP=PAR})]$ (Ar = Ph **18**, *para*-Tol **19**) were identified as being the third component upon subsequent preparation using more forcing conditions (vide infra).




Scheme 7



Scheme 8

Increasing the phosphine bulk proved one way to direct the reactivity of **A**. Thus treatment with mesitylphosphine (PMesH_2) gave ca. 95% formation (based on NMR

spectroscopy) of only the phosphido hydrido complex, *trans*-[Ni(I^tPr_2)₂(PMeSH)H] (**20**, Scheme 9). As for **10** and **12**, this was readily characterised from the ^1H NMR spectrum through the presence of a low frequency Ni-H resonance (ca. $\delta = -11$ ppm) with a $^2J(\text{H,P})$ doublet splitting (61 Hz) again indicative of the hydride being *trans* to the phosphido ligand. An additional smaller coupling of 2 Hz to the PMeSH proton was observed. The P-H resonance itself appeared at $\delta = 2.9$ ppm with a doublet $^1J(\text{H,P})$ coupling of 200 Hz; as noted by others, the magnitude of this splitting is similar to that in the free primary phosphine.^[39] In the proton coupled ^{31}P NMR spectrum, the phosphido ligand appeared as a very low frequency doublet of doublets ($\delta = -91$ ppm), while the $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum displayed the high frequency ($\delta = 191$ ppm) carbenic carbon resonance as the anticipated doublet with a *cis*-NHC-Ni-PR₂ $^2J(\text{C,P})$ coupling of 9 Hz.



Scheme 9

Given the similarity of the NMR spectra of **20** to those of **10**, **12**, **14** and **15**, it is reasonable to assume that these four compounds also displayed *trans* H-Ni-PArH geometries; they therefore provide a direct comparison of spectroscopic data for isostructural primary and secondary phosphido hydrido complexes. Thus, the Ni-H resonances of the former resonated at slightly higher frequency and showed smaller $^2J(\text{H,P})$ coupling constants (**10**: $\delta = -10.04$

ppm, $^2J(\text{H,P}) = 58.0$ Hz c.f. **5**: $\delta = -10.97$ ppm, $^2J(\text{H,P}) = 68.0$ Hz; **14**: $\delta = -10.41$ ppm, $^2J(\text{H,P}) = 62.1$ Hz c.f. **3**: $\delta = -11.31$ ppm, $^2J(\text{H,P}) = 70.0$ Hz), while the ^{31}P NMR signals appeared at significantly lower frequencies (**10**: $\delta = -43.8$ ppm; **12**: $\delta = -38.1$ ppm; **14**: $\delta = -41.2$ ppm c.f. **5**: $\delta = 19.3$ ppm; **4**: $\delta = 18.1$ ppm; **3**: $\delta = 25.1$ ppm; c.f. Table 1).^[40]

Characterisation of *trans*-[Ni(NHC)₂(PArH)₂] species. The bis-phosphido complexes **11** and **13** (Scheme 7) formed as the ultimate products of the reactions of **B** and **C** with PPhH₂ over several days at room temperature. Similarly, despite the bulk of PMesH₂, **20** also slowly converted to *trans*-[Ni(*i*Pr₂)₂(PMesH)₂] (**21**, Scheme 9) upon addition of further PMesH₂, albeit in only low yield. At the opposite end of the steric spectrum to **21** was *trans*-[Ni(*i*Pr₂)₂(PH₂)₂] (**22**), which formed as the major product of the reaction of **A** with excess PH₃ (Scheme 9).^[41]

Complexes **11**, **16**, **17**, **21** and **22** were structurally characterized as shown in Figure 4. They represent not only rare cases of structurally characterized metal primary phosphido complexes^[21c,40,42] but are the first examples of *trans*-RHP-M-PRH complexes. Table 1 summarises selected solid-state and spectroscopic metrics for these compounds, together with the two bis-PPh₂ complexes **6** and **7** for comparison. In general, the Ni-PArH distances are slightly shorter than those in the Ni-PAr₂ species (with the exception of the sterically crowded bis-P(Mes)H complex **21**). Although the *trans* PR₂-Ni-PR₂ geometry may limit comparisons to other systems, Glueck has reported that the Pd-PR₂ distance in [Pd(dppe)(PMes₂)Me] > [Pd(dppe)(P(Mes)H)Me] and that the *trans* influence of PPh₂ > PHPh for the [Pd(dppe)(PR₂)Me] complexes.^[39] The bis-PH₂ complex **22** is notable in displaying significant upfield shifts of both the ^1H and ^{31}P signals of PH₂ ligands, as well as a much reduced $^1J(\text{H,P})$ coupling constant.^[43]








Figure 4. Molecular structures (from top left to bottom right) of *trans*-[Ni(IME₄)₂(PPhH)₂] (**11**; ⁱ 1 – x, 1 – y, 1 – z symmetry operation), *trans*-[Ni(IⁱPr₂)₂(PPhH)₂] (**16**), *trans*-[Ni(IⁱPr₂)₂(P{*para*-Tol)H}₂)] (**17**; ⁱ 1 – x, 1 – y, – z symmetry operation), *trans*-[Ni(IⁱPr₂)₂(PMesH)₂] (**21**) and *trans*-[Ni(IⁱPr₂)₂(PH₂)₂] (**22**; ⁱ 1 – x, 2 – y, 1 – z symmetry operation). Ellipsoids are represented at 30% probability and hydrogens (with the exception of those bonded to phosphorus) have been omitted for clarity. Also omitted is the minor disordered component in **16** and lattice solvent in **21**. Only one of the two molecules in the asymmetric unit of **22** is shown.

	6	7	11	16	17	21	22
r(Ni-P)	2.2628(5)	2.2668(3)	2.2349(3)	2.2399(13)	2.2396(13)	2.2671(8)	2.2241(9)
				2.2414(13)		2.2550(8)	
r(Ni-C _{NHC})	1.8893(18)	1.8905(12)	1.8917(12)	1.896(4)	1.8852(19)	1.885(3)	1.890(3)
				1.888(4)		1.899(3)	
∠(P-Ni-P)	179.999(16)	180.00	180.00	175.42(6)	180.000(17)	172.12(3)	180.00
Σ(angles at P)	327.7	329.6	307.1	311.7/313.2	314.2	320.2/321.3	306.8
δ ¹ H _{PH}	-	-	3.33	3.11	3.11	2.94	0.90
³¹ P	31.2	47.1	-40.4	-29.5	-31.0	-71.2	-156.3
¹ J(H,P)	-	-	-	198	198	200	171
δ ¹³ C _{NHC}	182.7	190.3	188.1	186.3	190.0	185.9	190.0
² J(C _{NHC} ,P)	16	13	13	14	13	13	11

Table 1. Comparison of bond distances [Å] and angles [deg] and selected NMR spectroscopic data (δ [ppm], *J* [Hz]) for the bis-phosphido complexes **6**, **7**, **11**, **16**, **17**, **21** and **22**.

Isolation and characterization of $[\text{Ni}(\text{I}^i\text{Pr}_2)_2(\eta^2\text{-ArP=PAr})]$. The diphosphene complexes $[\text{Ni}(\text{I}^i\text{Pr}_2)_2(\eta^2\text{-ArP=PAr})]$ (Ar = Ph **18**, *para*-Tol **19**) were isolated in ca. 50% yield by heating **A** with excess PPhH_2 or $\text{P}(\text{para-Tol})\text{H}_2$ at 140 °C. While diphosphene complexes species have been synthesised previously using a range of approaches (ligand exchange of stable diphosphenes with labile M-L complexes, salt metathesis and degradation of oligophosphines),^[30e, 44] their preparation via the formal dehydrocoupling of a primary phosphine in the coordination sphere of a transition metal as observed here is unusual.^[45]

X-ray crystallography (Figure 5) showed, in both cases, a *trans* disposition of the two aryl substituents in the diphosphene ligand. The P-P bond lengths (**18**: 2.1379(7) Å; **19**: 2.1342(7) Å) were in good agreement with those in $[\text{Ni}(\text{PEt}_3)_2(\text{trans-MesP=PMes})]$ (2.1355(9) Å), $[\text{Ni}(\text{PMe}_2\text{Ph})_2(\text{trans-MesP=PMes})]$ (2.137(1) Å), and $[\text{Pd}(\text{dppe})(\text{trans-PhP=PPh})]$ (2.121(4) Å; dppe = 1,2-bis(diphenylphosphino)ethane), respectively.^[44,46] The angles C1–Ni–C11 (**18**, 104.62(6)°; **19**, 98.85(8)°) were larger than 90°, as expected for a pseudo three-coordinated complex with small P-Ni-P angles (**18**, 56.76(2)°; **19**, 56.89(2)°).

The ^1H NMR resonances of the I^iPr_2 ligands in both **18** and **19** were broadened at room temperature, consistent with hindered rotation of the ArP=PAr ligand around the Ni-(PP)_{centroid} vector and/or of the NHC ligands around the Ni-C bonds. Cooling to -50 °C resolved the spectra into four ^iPr methyl and two ^iPr methine resonances, as typically found for a C_{2v} type structure. At 90 °C, two ^iPr methyl resonances and a single ^iPr methine signal were observed, which were still broadened. Both complexes exhibited singlets in their $^{31}\text{P}\{^1\text{H}\}$ NMR spectra (**18**, $\delta = -41$ ppm; **19**, $\delta = -39$ ppm) to lower frequency of the Pd and Pt derivatives $[\text{M}(\text{dppe})(\text{trans-PhP=PPh})]$ (M = Pd, $\delta = -34$ ppm; M = Pt, $\delta = -24$ ppm).^[44]




Figure 5. Molecular structures of $[\text{Ni}(\text{I}'\text{Pr})_2(\eta^2\text{-PhP=PPh})]$ (**18**) and $[\text{Ni}(\text{I}'\text{Pr})_2\{\eta^2\text{-(para-Tol)P=P(para-Tol)}\}]$ (**19**). Ellipsoids are represented at 30% probability and hydrogens (along with lattice solvent in **19**) have been omitted for clarity. For **19**: $i\ 1 - x, y, \frac{1}{2} - z$ symmetry operation.

Computational Studies. (i) P-H Activation of PPh_2H at $[\text{Ni}(\text{NHC})_2]$. The mechanisms of these processes were investigated using DFT calculations for the reactions of both $[\text{Ni}(\text{I}'\text{Pr})_2]$, **A**, and $[\text{Ni}(\text{IME}_4)_2]$, **B**. Geometry optimisation and frequency calculations were performed with the BP86 functional using SDD pseudopotentials and basis sets on Ni and P (with d-orbital polarization on the latter) and 6-31G** basis sets on all other atoms (collectively termed BS1). Electronic energies were then recomputed with B97D including corrections for benzene solvent and a def2-TZVP basis set (BS2). These were then combined with the BP86 thermochemical corrections to supply the relative free energies reported in the text. The choice of B97D was motivated by the computed barriers for the reactions of **A** and **B** with PPh_2H , for which this functional gave values qualitatively consistent with the reactivities observed at room temperature (i.e. **B**: rapid formation of *trans*- $[\text{Ni}(\text{IME}_4)_2(\text{PPh}_2)\text{H}]$, **5**, followed by slower conversion to *trans*- $[\text{Ni}(\text{IME}_4)_2(\text{PPh}_2)_2]$, **6**; **A**: rapid formation of *trans*- $[\text{Ni}(\text{I}'\text{Pr})_2(\text{PPh}_2)\text{H}]$, **3**, but no onward reaction to form *bis*-phosphido complex **8**). The

computed energetics were found to be very functional dependent, in a similar way that phosphine ligand addition energies are highly sensitive to functional choice and, in particular, the treatment of dispersion effects.^[47,48] This appears to be exacerbated here due to the sequential addition of two PPh₂H ligands to two [Ni(NHC)₂] systems of varying steric bulk. The data below therefore present our best proposed mechanisms in the face of these challenges.^[49] Full computational details, references and functional testing are provided in the Supporting Information.

The free energy profile computed with B97D for the reaction of **B** with PPh₂H to form **5** is shown in Figure 6. Addition of PPh₂H to **B** is exergonic by 11.6 kcal/mol and forms [Ni(Ime₄)₂(PPh₂H)], **I**, the energy of which is set to 0.0 kcal/mol.^[50] P-H activation then proceeds via an initial rearrangement to intermediate **II**, in which the H substituent bridges the Ni-P bond (Ni-H = 1.62 Å, P-H = 1.59 Å, Ni-P = 2.18 Å). The P-H bond is therefore significantly weakened and readily undergoes cleavage to form the *cis* isomer of **5**, denoted *cis*-**III** in the computational study. Formation of *cis*-**III** is slightly endergonic, and hence reversible, allowing the *trans* isomer to be accessed via **TS(II-*trans*-III)** at +19.6 kcal/mol (see Figure 7 for the computed geometries of rate-limiting transition states). The much higher barrier associated with **TS(II-*trans*-III)** arises from the coupling of P-H bond cleavage with a *cis-trans* isomerisation process. The triplet form of [Ni(Ime₄)₂(PPh₂H)], ³**III**, has a computed energy of +23.7 kcal/mol and so is not implicated in this isomerisation.^[51] Ime₄ dissociation from *cis*-**III** was also ruled out as this incurs a free energy penalty of 25.7 kcal/mol. Overall, *trans*-**III** is formed with an overall barrier of 19.6 kcal/mol in a process that is exergonic by 2.4 kcal/mol relative to **I**.





Figure 6. Computed free energy profile [kcal/mol, B97D(benzene, BS2)//BP86(BS1)] for the reaction of [Ni(Ime₄)₂] (**B**) with PPh₂H to form *trans*-[Ni(Ime₄)₂(PPh₂)H] **5**. The triplet form of [Ni(Ime₄)₂(PPh₂)H], **³III**, is computed at +23.7 kcal/mol.

The lowest energy pathway for the onward reaction of **5** with a second molecule of PPh₂H to form **6** is shown in Figure 8.^[52] This involves an initial P-H reductive coupling to reform **II** which can then add a second PPh₂H molecule in the unconventional H-bridged binding mode to give **IV** (+10.7 kcal/mol). Both P-H bonds in **IV** are significantly elongated (ca. 1.63 Å) and a P-H activation transition state, **TS(IV-V)**, was located at +21.9 kcal/mol in which both P-H bonds are cleaved (see Figure 8). This occurs with concomitant H₂ elimination to give *cis*-**V** directly.^[53] *cis*-**V** can then isomerise to *trans*-**V** (i.e. **6**) at -19.9 kcal/mol via **TS(*cis-trans*-V)** at +4.9 kcal/mol on the singlet surface, although in this case the triplet species **³V** lies 3.8 kcal/mol below **TS(*cis-trans*-V)** and so isomerisation may involve a spin crossover process.^[54] These isomerisation pathways do not affect the overall energy span of 24.3 kcal/mol for the second P-H bond activation, as this is associated with **TS(IV-V)** as the highest point on the profile. The formation of **6** from **5** is strongly exergonic (-17.5 kcal/mol).

1st P–H Activation



2nd P–H Activation




Figure 7. Computed geometries for the rate-limiting transition states for the first and second P-H activation of PPh_2H at **B**. Selected distances are shown in Å, with IMe_4 and phenyl hydrogens omitted for clarity.




Figure 8. Computed free energy profile [kcal/mol, B97D(benzene, BS2)//BP86(BS1)] for the reaction of **5** with PPh_2H to form *trans*- $[\text{Ni}(\text{IMe}_4)_2(\text{PPh}_2)_2]$ **6**.

Overall, the first P-H activation at **B** to form **5** is more kinetically accessible ($\Delta G^\ddagger = 19.6$ kcal/mol) than the second to form **6** ($\Delta G^\ddagger = +24.3$ kcal/mol) and this is qualitatively consistent with the rapid formation of **5** at room temperature and its subsequent slower conversion to **6**. Calculations for the reactions of **A** with PPh₂H followed similar pathways, with a barrier of 20.2 kcal/mol for the first P-H activation to form **3** at -0.8 kcal/mol. In this case the potential second P-H bond activation entails a higher overall barrier of 26.6 kcal/mol, consistent with the formation of *trans*-[Ni(I^{*i*}Pr)₂(PPh₂)₂], **8**, being much less accessible at room temperature, at least via this route (see Figures S74 and S75 in the Supporting Information).^[55]

As well as the observed P-H bond activations, the alternative P-Ph bond activation of PPh₂H at [Ni(NHC)]₂ is a potentially competing process (see Figure 9 for the IMe₄ system). This proceeds with an initial computed barrier of 13.4 kcal/mol to give *cis*-[Ni(IMe₄)₂(PPhH)Ph], *cis*-**III**_{Ph}, at -1.6 kcal/mol from which *cis-trans* isomerisation entails a barrier of 25.6 kcal/mol to give *trans*-**III**_{Ph} at -6.2 kcal/mol. Therefore, in the absence of a second equivalent of PPh₂H, P-H bond activation to form **6** is kinetically favoured over *trans*-**III**_{Ph} ($\Delta G^\ddagger = +19.6$ kcal/mol, $\Delta G = -2.4$ kcal/mol) but as this could be reversible, this could in principle allow access to *trans*-**III**_{Ph} at -6.2 kcal/mol as the thermodynamically more stable product of P-C bond activation. This latter process, however, would have a significant barrier of 26.4 kcal/mol relative to **6**. Moreover, in the presence of excess PPh₂H, a second P-H activation to form **6** is favoured both kinetically ($\Delta G^\ddagger = +24.3$ kcal/mol) and thermodynamically ($\Delta G = -19.9$ kcal/mol). The pattern of a kinetically favoured P-H bond activation vs. a thermodynamically favoured P-C bond activation has been reported for the reaction of Pt(dppe)(*trans*-stilbene) with PMes₂H.^[56,57] Attempts to realise this pattern experimentally through the reaction of Ni(COD)₂ with 2 equiv of IMe₄ and only one equiv of PPh₂H were unsuccessful with a complex mixture of products being formed.




Figure 9. Computed free energy profile [kcal/mol, B97D(benzene, BS2)//BP86(BS1)] for competing P-C bond activation in $[\text{Ni}(\text{IME}_4)_2(\text{PPh}_2\text{H})]$, **I**. The triplet form of $[\text{Ni}(\text{IME}_4)_2(\text{PPh}_2\text{Ph})]$, $^3\text{IV}_{\text{Ph}}$, is computed at +25.6 kcal/mol.

(ii) P-P Activation of $\text{Ph}_2\text{P-PPh}_2$ at $[\text{Ni}(\text{NHC})_2]$. The computed profile for the alternative route to $\text{trans-}[\text{Ni}(\text{IME}_4)_2(\text{PPh}_2)_2]$, **6**, via reaction of $\text{Ph}_2\text{P-PPh}_2$ with **B** is shown in Figure 10 with geometries of key transition states in Figure 11. $\text{Ph}_2\text{P-PPh}_2$ addition to **B** is exergonic by 17.8 kcal/mol (*cf.* 11.6 kcal/mol for PPh_2H) and gives trigonal planar $[\text{Ni}(\text{IME}_4)_2(\text{Ph}_2\text{PPPh}_2)]$ (**VI**, Ni-P = 2.35 Å), the energy of which is set to 0.0 kcal/mol. P-P bond activation then proceeds via a facile transfer of one PPh_2 group onto Ni via **TS(VI-*cis*-V)** at +9.0 kcal/mol. This forms *cis*-**V** at -5.5 kcal/mol that, as discussed above, can undergo *cis-trans* isomerisation on the singlet surface via **TS(*cis-trans*-VI)** at +10.3 kcal/mol or potentially via a lower energy spin crossover mechanism via ^3V at +6.5 kcal/mol.^[58] *Cis-trans* isomerisation on the singlet surface represents the largest barrier along the profile for the formation of **6**, but at only 15.8 kcal/mol, the implication is that the overall reaction to form **6** should be readily accessible, consistent with the experiment observation of this process at room temperature.




Figure 10. Computed free energy profiles [kcal/mol, B97D(benzene, BS2)/BP86(BS1)] for the reaction of **B** with $\text{Ph}_2\text{P-PPh}_2$ to form **6**.




Figure 11. Computed geometries for key transition states in the P-P bond activation of Ph_2PPPh_2 at **B**. Selected distances are shown in Å, with IMe_4 and phenyl hydrogens omitted for clarity.

P-P bond activation in the analogous reaction of $\text{Ph}_2\text{P-PPh}_2$ with **A** is shown in Figure 12. This now involves a two-step process from the initial $\text{Ph}_2\text{P-PPh}_2$ adduct, VI_{iPr} , via an intermediate $\text{Int}(\text{VI-}i\text{cis-V})_{\text{iPr}}$ at -0.3 kcal/mol that features a bridging PPh_2 ligand, followed by facile P-P bond cleavage to give $i\text{cis-V}_{\text{iPr}}$ at -5.8 kcal/mol. *Cis-trans* isomerisation on the singlet surface then proceeds via $\text{TS}(i\text{cis-trans-V})_{\text{iPr}}$ at +11.8 kcal/mol. The details of this process are complicated by the existence of numerous conformations along the pathway and only the highest-lying transition state is shown in Figure 12 (see Figure S78 in the Supporting Information for more details). The triplet form, $^3\text{V}_{\text{iPr}}$, is again accessible at +4.3 kcal/mol. As with the analogous reaction with **B**, all barriers are readily accessible, the largest energy span being only 17.6 kcal/mol (corresponding to isomerisation on the singlet surface) and so the formation of **8** should be readily accessible, as is the case experimentally.




Figure 12. Computed free energy reaction profile [kcal/mol, B97D(benzene, BS2)//BP86(BS1)] for the reaction of **A** with $\text{Ph}_2\text{P-PPh}_2$ to form *trans*-[Ni(*i*Pr)₂(PPh₂)₂], **8**; ^a only the rate-limiting transition state for *cis-trans* isomerisation is indicated (see Figure S78 in the Supporting Information for more details).


Summary and Conclusions.

Facile P-H activation of secondary and primary phosphines by the 14-electron fragments $[\text{Ni}(\text{I}^i\text{Pr}_2)_2]$ (**A**), $[\text{Ni}(\text{IMe}_4)_2]$ (**B**) and $[\text{Ni}(\text{IEt}_2\text{Me}_2)_2]$ (**C**) provides access to a unprecedented series of stable terminal phosphido complexes with *trans* H-Ni-PR₂ and *trans*-R₂P-Ni-PR₂ structures. Within the family of $[\text{Ni}(\text{NHC})_2]$ fragments that we have considered, clear evidence for the stereoelectronic influence of the N-substituents is apparent. This is best evidenced in the initial reactions of **A**, **B** and **C** with PPh₂H. With bulkier **A** and **C**, the initial room temperature reaction yields isolable phosphido hydrido complexes, *trans*- $[\text{Ni}(\text{I}^i\text{Pr}_2)_2(\text{PPh}_2)\text{H}]$ (**3**) and *trans*- $[\text{Ni}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_2)\text{H}]$ (**4**), respectively. With **B**, a mixture of the equivalent phosphido hydrido *trans*- $[\text{Ni}(\text{IMe}_4)_2(\text{PPh}_2)\text{H}]$ (**5**) and bis-phosphido *trans*- $[\text{Ni}(\text{IMe}_4)_2(\text{PPh}_2)_2]$ (**6**) complexes is first seen that then converts fully to **6** over 24 h. Thus **B**, that features the smallest of the carbenes, IMe₄ can most easily accommodate two P-H bond activations to form a *bis*-phosphido directly. Further reaction of **4** with PPh₂H to form *trans*- $[\text{Ni}(\text{IEt}_2\text{Me}_2)_2(\text{PPh}_2)_2]$ (**7**) does occur but is very slow, requiring weeks at room temperature. No further reaction between **3** and PPh₂H is seen and so $[\text{Ni}(\text{I}^i\text{Pr}_2)_2(\text{PPh}_2)_2]$ (**8**) is inaccessible by this route. We were able to prepare **8**, as well as **7** and the mixed aryl/alkyl bis-phosphido complex $[\text{Ni}(\text{I}^i\text{Pr}_2)_2(\text{PPhMe}_2)_2]$ (**9**) by the unexpected P-P cleavage of the diphosphines R₂P-PR₂, which like the P-H activation reactions above, takes place under very mild conditions.

These trends were captured by DFT calculations on the reactions of **A** and **B**, although the computed energetics of these processes were found to be very functional dependent. With PPh₂H low barriers for the first P-H activation (**A**: 20.2 kcal/mol; **B**: 19.6 kcal/mol, B97D functional) contrasted with increased barriers for the second P-H activation (**A**: 26.6 kcal/mol; **B**: 24.3 kcal/mol). Each P-H bond activation proceeds from an unconventional phosphine adduct in which the H substituent bridges the Ni-P bond. The rate limiting transition states then correspond to P-H bond cleavage; for the first P-H activation, this is coupled to *cis-trans* isomerisation to form *trans*- $[\text{Ni}(\text{NHC})_2(\text{PPh}_2)(\text{H})]$, whereas for the second P-H activation, a novel transition state involving dual P-H bond cleavage of two

PPh₂H ligands is located. P-P bond activation from adducts of the type [Ni(NHC)₂(Ph₂P-PPh₂)] (NHC = IⁱPr, IMe₄,) is computed to be an extremely facile process, with barriers below 10 kcal/mol to form *cis*-[Ni(NHC)₂(PPh₂)₂]. In this case *cis-trans* isomerisation has a higher barrier but remains readily accessible.

The ability of the [Ni(NHC)₂] moiety to afford isolable terminal phosphido products is in marked contrast to previous efforts to prepare analogues based on a [Ni(PR₃)₂] core, all of which have been reported to be unstable even at low temperatures. The latter often react on to form bridging phosphido dimers and higher nuclearity species,^[20] suggesting phosphine loss as a key process in this onward reactivity. The higher binding energies of NHC ligands^[59] may therefore contribute to the greater stabilities exhibited here. To quantify this hypothesis, calculations on ligand dissociation and subsequent dimer formation were performed on *trans*-[NiL₂(PH₂)₂] (L = PH₃, IMe₂; IMe₂ = 1,3-dimethylimidazol-2-ylidene) where these simple model systems were employed to probe electronic effects directly.



Scheme 10. Computed free energies [kcal/mol, B97D(benzene, BS2)//BP86(BS1)] for ligand loss and dimerization in *trans*-[NiL₂(PH₂)₂] (L = PH₃, IMe₂).

As shown in Scheme 10, phosphine dissociation from *trans*-[Ni(PH₃)₂(PH₂)₂] is significantly more accessible than NHC loss from *trans*-[Ni(IMe₂)₂(PH₂)₂]. Moreover, subsequent dimer formation is exergonic for the phosphine system, but strongly endergonic with the NHC analogue. This provides some insight into the remarkable ability of the

[Ni(NHC)₂] fragment to not only effect a wide range of novel bond activation processes, but also to stabilize the resultant products.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

We thank the Royal Society (Newton Fellowship to SS), EPSRC, the Deutsche Forschungsgemeinschaft (DFG; RA720/7), the Julius-Maximilians-Universität Würzburg and Heriot-Watt University for financial support.

Keywords

carbene ligands

density functional calculations

hydride ligands

nickel

phosphido ligands

References

- [1] A. J. Arduengo, S. F. Gamper, J. C. Calabrese, F. Davidson, *J. Am. Chem. Soc.* **1994**, *116*, 4391-4394.
- [2] a) K. Matsubara, S. Miyazaki, Y. Koga, Y. Nibu, T. Hashimura, T. Matsumoto, *Organometallics* **2008**, *27*, 6020-6024; b) A. A. Danopoulos, D. Pugh, *Dalton Trans.* **2008**, 30-31; c) R. C. Poulten, M. J. Page, A. G. Algarra, J. J. Le Roy, I. López, E. Carter, A. Llobet, S. A. Macgregor, M. F. Mahon, D. M. Murphy, M. Murugesu, M. K. Whittlesey, *J. Am. Chem. Soc.* **2013**, *135*, 13640-13643; d) Y. Hoshimoto, Y. Hayashi, H. Suzuki, M. Ohashi, S. Ogoshi, *Organometallics* **2014**, *33*, 1276-1282.
- [3] For an example with a cyclic (alkyl)(amino)(carbene) in which the oxidation state lies between Ni(0) and Ni(I), see: K. C. Mondal, P. P. Samuel, Y. Li, H. W. Roesky, S. Roy, L. Ackermann, N. S. Sidhu, G. M. Sheldrick, E. Carl, S. Demeshko, S. De, P. Parameswaran, L. Ungur, L. F. Chibotaru, D. M. Andrada, *Eur. J. Inorg. Chem.* **2014**, 818-823.
- [4] a) N. D. Clement, K. J. Cavell, C. Jones, C. J. Elsevier, *Angew. Chem. Int. Ed.* **2004**, *43*, 1277-1279; *Angew. Chem.* **2004**, *116*, 1297-1299; b) N. D. Clement, K. J. Cavell, L. Ooi, *Organometallics* **2006**, *25*, 4155-4165; c) S. Miyazaki, Y. Koga, T. Matsumoto, K. Matsubara, *Chem. Commun.* **2010**, *46*, 1932-1934; d) K. Zhang, M. Conda-Sheridan, S. R. Cooke, J. Louie, *Organometallics* **2011**, *30*, 2546-2552; e) C. H. Lee, T. R. Cook, D. G. Nocera, *Inorg. Chem.* **2011**, *50*, 714-716; f) N. D. Harrold, A. R. Corcos, G. L. Hillhouse, *J. Organomet. Chem.* **2016**, *813*, 46-54; g) D. J. Nelson, F. Maseras, *Chem. Commun.* **2018**, *54*, 10646-10649; h) L. Tendra, T. Schaub, M. J. Krahfuss, M. W. Kuntze-Fechner, U. Radius, *Eur. J. Inorg. Chem.* **2020**, 3194-3207.

- [5] The high reactivity of [Ni(IMes)₂] towards C-F bonds of fluoroarenes and a comparison with the reactivity of [Ni₂(I^{*i*}Pr)₄(COD)] in C-F bond activation has very recently been reported. a) M. W. Kuntze-Fechner, H. Verplancke, L. Tendera, M. Diefenbach, I. Krummenacher, H. Braunschweig, T. B. Marder, M. C. Holthausen, U. Radius, *Chem. Sci.* **2020**, *11*, 11009–11023. For reports of C-F, C-Cl, and C-H borylation using [Ni(IMes)₂] as a catalyst see; b) J. Zhou, M. Kuntze-Fechner, R. Bertermann, U. S. D. Paul, J. H. J. Berthel, A. Friedrich, Z. Du, T. B. Marder, U. Radius, *J. Am. Chem. Soc.* **2016**, *138*, 5250-5253; c) Y. Tian, X. Guo, M. Kuntze-Fechner, I. Krummenacher, H. Braunschweig, U. Radius, A. Steffen, T. B. Marder, *J. Am. Chem. Soc.* **2018**, *140*, 17612-17623; d) Y. Tian, X. Guo, Z. Wu, A. Friedrich, H. Braunschweig, U. Radius, T. B. Marder, *J. Am. Chem. Soc.* **2020**, *142*, 13136-13144; e) Y. Tian, X. Guo, I. Krummenacher, Z. Wu, J. Nitsch, H. Braunschweig, U. Radius, T. B. Marder, *J. Am. Chem. Soc.* **2020**, *142*, 18231-18242.
- [6] For a recent overview, see: A. A. Danopoulos, T. Simler, P. Braunstein, *Chem. Rev.* **2019**, *119*, 3730-3961.
- [7] P. L. Arnold, F. G. N. Cloke, T. Geldbach, P. B. Hitchcock, *Organometallics* **1999**, *18*, 3228-3233.
- [8] a) T. Schaub, U. Radius, *Chem. Eur. J.* **2005**, *11*, 5024-5030; b) T. Schaub, M. Backes, U. Radius, *Organometallics* **2006**, *25*, 4196-4206.
- [9] a) P. Fischer, T. Linder, U. Radius, *Z. Anorg. Allg. Chem.* **2012**, *638*, 1491-1496; b) J. H. J. Berthel, L. Tendera, M. W. Kuntze-Fechner, L. Kuehn, U. Radius, *Eur. J. Inorg. Chem.* **2019**, 3061-3072; c) L. Kuehn, D. G. Jammal, K. Lubitz, T. B. Marder, U. Radius, *Chem. Eur. J.* **2019**, *25*, 9514-9521.
- [10] a) T. Schaub, P. Fischer, A. Steffen, T. Braun, U. Radius, A. Mix, *J. Am. Chem. Soc.* **2008**, *130*, 9304-9317; b) T. Schaub, M. Backes, U. Radius, *Eur. J. Inorg. Chem.*

- 2008**, 2680-2690; c) T. Schaub, P. Fischer, T. Meins, U. Radius, *Eur. J. Inorg. Chem.* **2011**, 3122-3126; d) H. Iwamoto, H. Imiya, M. Ohashi, S. Ogoshi, *J. Am. Chem. Soc.* **2020**, *142*, 19360-19367.
- [11] a) T. Zell, M. Feierabend, B. Halfter, U. Radius, *J. Organomet. Chem.* **2011**, *696*, 1380-1387; b) T. Zell, P. Fischer, D. Schmidt, U. Radius, *Organometallics* **2012**, *31*, 5065-5073; c) T. Zell, U. Radius, *Z. Anorg. Allg. Chem.* **2013**, *639*, 334-339.
- [12] a) T. Schaub, M. Backes, U. Radius, *Chem. Commun.* **2007**, 2037-2039; b) T. Schaub, M. Backes, O. Plietzsch, U. Radius, *Dalton Trans.* **2009**, 7071-7079.
- [13] a) T. Zell, T. Schaub, K. Radacki, U. Radius, *Dalton Trans.* **2011**, *40*, 1852-1854; b) C. Hauf, J. E. Barquera-Lozada, P. Meixner, G. Eickerling, S. Altmannshofer, D. Stalke, T. Zell, D. Schmidt, U. Radius, W. Scherer, *Z. Anorg. Allg. Chem.* **2013**, *639*, 1996-2004.
- [14] a) T. Schaub, M. Backes, U. Radius, *J. Am. Chem. Soc.* **2006**, *128*, 15964-15965; b) J. Zhou, J. H. J. Berthel, M. W. Kuntze-Fechner, A. Friedrich, T. B. Marder, U. Radius, *J. Org. Chem.* **2016**, *81*, 5789-5794; c) M. W. Kuntze-Fechner, C. Kerpen, D. Schmidt, M. Häring, U. Radius, *Eur. J. Inorg. Chem.* **2019**, 1767-1775.
- [15] P. Fischer, K. Götz, A. Eichhorn, U. Radius, *Organometallics* **2012**, *31*, 1374-1383.
- [16] D. Schmidt, T. Zell, T. Schaub, U. Radius, *Dalton Trans.* **2014**, *43*, 10816-10827.
- [17] T. Schaub, C. Döring, U. Radius, *Dalton Trans.* **2007**, 1993-2002.
- [18] S. Sabater, M. J. Page, M. F. Mahon, M. K. Whittlesey, *Organometallics* **2017**, *36*, 1776-1783.
- [19] a) J. Chatt, J. M. Davidson, *J. Chem. Soc.* **1964**, 2433-2445; b) R. Bender, P. Braunstein, A. Dedieu, P. D. Ellis, B. Huggins, P. D. Harvey, E. Sappa, A. Tiripicchio, *Inorg. Chem.* **1996**, *35*, 1223-1234; c) C. Scriban, D. K. Wicht, D. S. Glueck, L. N. Zakharov, J. A. Golen, A. L. Rheingold, *Organometallics* **2006**, *25*, 3370-3378; d) P.

- Mastrorilli, *Eur. J. Inorg. Chem.* **2008**, 4835-4850; e) R. Beck, M. Shoshani, J. Krasinkiewicz, J. A. Hatnean, S. A. Johnson, *Dalton Trans.* **2013**, 42, 1461-1475; f) V. Varela-Izquierdo, A. M. Geer, B. de Bruin, J. A. López, M. A. Ciriano, C. Tejel, *Chem. Eur. J.* **2019**, 25, 15915-15928; g) A. Berkefeld, M. Reimann, G. Hörner, M. Kaupp, H. Schubert, *Organometallics* **2020**, 39, 443-452.
- [20] D. R. Fahey, J. E. Mahan, *J. Am. Chem. Soc.* **1976**, 98, 4499-4503.
- [21] a) H. Schäfer, *Z. Anorg. Allg. Chem.* **1979**, 459, 157-169; b) H. Schäfer, D. Binder, B. Deppisch, G. Mattern, *Z. Anorg. Allg. Chem.* **1987**, 546, 79-98; c) R. Melenkivitz, D. J. Mindiola, G. L. Hillhouse, *J. Am. Chem. Soc.* **2002**, 124, 3846-3847; d) M. Madadi, B. K. Najafabadi, M. A. Fard, J. F. Corrigan, *Eur. J. Inorg. Chem.* **2015**, 3094-3101; e) J. A. Kessler, V. M. Iluc, *Dalton Trans.* **2017**, 46, 12125-12131.
- [22] For very recent examples, see: a) T. J. Hadlington, T. Szilvási, M. Driess, *J. Am. Chem. Soc.* **2019**, 141, 3304-3314; b) K. Lee, C. E. Moore, C. M. Thomas, *Organometallics* **2020**, 39, 2053-2056.
- [23] a) Y. S. Ganushevich, V. A. Miluykov, F. M. Polyancev, S. K. Latypov, P. Loennecke, E. Hey-Hawkins, D. G. Yakhvarov, O. G. Sinyashin, *Organometallics* **2013**, 32, 3914-3919; b) S. K. Latypov, F. M. Polyancev, Y. S. Ganushevich, V. A. Miluykov, O. G. Sinyashin, *Dalton Trans.* **2016**, 45, 2053-2059.
- [24] Deposition Number(s) <https://www.ccdc.cam.ac.uk/services/structures?id=doi:10.1002/chem.20210XXXX> 2009924 (for **2**), 2009925 (for **3**), 2009926 (for **4**), 2009927 (for **6**), 2009928 (for **7**), 2009929 (for **11**), 2009930 (for **16**), 2009931 (for **17**), 2009932 (for **18**), 2009933 (for **19**), 2009934 (for **21**), 2009935 (for **22**), 2009936 (for $[\text{Ni}(\text{IEt}_2\text{Me}_2)]_2(\mu\text{-PPh}_2)_2$, Supporting Information only) contain(s) the supplementary crystallographic data for this paper. These data are provided free of

charge by the joint Cambridge Crystallographic Data Centre and

Fachinformationszentrum Karlsruhe <url href="

<http://www.ccdc.cam.ac.uk/structures>>Access Structures service</url>.

- [25] a) *Ir*: R. A. Schunn, *Inorg. Chem.* **1973**, *12*, 1573-1579; b) *Ir*: E. A. V. Ebsworth, R. A. Mayo, *Angew. Chem. Int. Ed.* **1985**, *24*, 68-70; *Angew. Chem.* **1985**, *97*, 65-66; c) *Ir*: E. A. V. Ebsworth, R. A. Mayo, *J. Chem. Soc., Dalton Trans.* **1988**, 477-484; d) *W*: L. D. Field, N. G. Jones, P. Turner, *J. Organomet. Chem.* **1998**, *571*, 195-199; e) *Ru*: V. S. Chan, I. C. Stewart, R. G. Bergman, F. D. Toste, *J. Am. Chem. Soc.* **2006**, *128*, 2786-2787.
- [26] This contrasts with *cis* phosphine-Ni-H splittings of ca. 65-77 Hz. M.Y. Darensbourg, M. Ludwig, C. G. Riordan, *Inorg. Chem.* **1989**, *28*, 1630-1634.
- [27] a) As reported in the Supporting Information, the phosphido hydrido complex **4** was crystallised from Et₂O/pentane and the bis-phosphido complex **7** from benzene/hexane. Small amounts of the green Ni(I) bridging-phosphido dimer, [Ni(IEt₂Me₂)]₂(μ-PPh₂)₂, could be isolated if either **4** or **7** were left in the presence of Et₂O for extended periods of time. This diamagnetic, dimeric compound was spectroscopically and structurally characterised (Supporting Information); b) For an analogue bearing P^{*i*}Pr₃ in place of the NHC, see ref [19e].
- [28] Even a single equivalent of PPh₂H still gave a mixture of **6** and **7**. Although this precluded isolation of **6**, NMR spectra (measured at low temperature to slow the conversion through to **7**) indicated it was isostructural with **3-5**.
- [29] Simply leaving **5** to stand in solution in the absence of any PPh₂H failed to give **10**, showing that bis-phosphido complex formation is not via two molecules of **5** combining, followed by ligand redistribution.

- [30] a) J. S. Figueroa, C. C. Cummins, *J. Am. Chem. Soc.* **2003**, *125*, 4020-4021; b) G. Bai, P. Wei, A. K. Das, D. W. Stephan, *Dalton Trans.* **2006**, 1141-1146; c) D. Yakhvarov, P. Barbaro, S. M. Carpio, S. Midollini, A. Orlandini, M. Peruzzini, O. Sinyashin, F. Zanobini, *Angew. Chem. Int. Ed.* **2006**, *45*, 4182-4185; *Angew. Chem.* **2006**, *118*, 4288-4291; d) S. Greenberg, D. W. Stephan, *Chem. Soc. Rev.* **2008**, *37*, 1482-1489; e) S. Gómez-Ruiz, S. Zahn, B. Kirchner, W. Böhlmann, E. Hey-Hawkins, *Chem. Eur. J.* **2008**, *14*, 8980-8985; f) S. J. Geier, D. W. Stephan, *Chem. Commun.* **2008**, 2779-2781; g) M. Arisawa, K. Sawahata, T. Yamad, D. Sarkar, M. Yamaguchi, *Org. Lett.* **2018**, *20*, 938-941; h) M. Modl, S. Heinl, G. Balázs, F. D. Calvo, M. Caporali, G. Manca, M. Keilwerth, K. Meyer, M. Peruzzini, M. Scheer, *Chem. Eur. J.* **2019**, *25*, 6300-6305.
- [31] a) S. J. Geier, D. W. Stephan, *Chem. Commun.* **2008**, 99-101; b) R. Waterman, *Curr. Org. Chem.* **2008**, *12*, 1322-1339.
- [32] T. Chu, Y. Boyko, I. Korobkov, L. G. Kuzmina, J. A. K. Howard, G. I. Nikonov, *Inorg. Chem.* **2016**, *55*, 9099-9104.
- [33] Activation of the P-P bond of an ortho-carborane-substituted 1,2-diphosphetane by the cobalt(-I) species $[K(thf)_{0.2}][Co(\eta^4-cod)_2]$ has been described by Hey-Hawkins, Wolf and co-workers. P. Coburger, S. Demeshko, C. Rödl, E. Hey-Hawkins, R. Wolf, *Angew. Chem. Int. Ed.* **2017**, *56*, 15871-15875; *Angew. Chem.* **2017**, *129*, 16087-16091.
- [34] For representative examples, see: a) V. P. W. Böhm, M. Brookhart, *Angew. Chem. Int. Ed.* **2001**, *40*, 4694-4696; *Angew. Chem.* **2011**, *113*, 4832-4834; b) L.-B. Han, T. D. Tilley, *J. Am. Chem. Soc.* **2006**, *128*, 13698-13699; c) A. M. Geer, Á. L. Serrano, B. de Bruin, M. A. Ciriano, C. Tejel, *Angew. Chem. Int. Ed.* **2015**, *54*, 472-475; *Angew. Chem.* **2015**, *127*, 482-485.

- [35] Structures of the homoleptic terminal phosphido species $[\text{Hf}(\text{PCy}_2)_5]^+$ and $[\text{Mo}(\text{PCy}_2)_4]$ have been reported. R. T. Baker, P. J. Krusic, T. H. Tulip, J. C. Calabrese, S. S. Wreford, *J. Am. Chem. Soc.* **1983**, *105*, 6763-6765.
- [36] For examples of bis-phosphido complexes with a *cis*- $[\text{R}_2\text{P}-\text{M}-\text{PR}_2]$ arrangement, see:
a) R. T. Baker, J. F. Whitney, S. S. Wreford, *Organometallics* **1983**, *2*, 1049-1051; b) L. Weber, G. Meine, R. Boese, N. Augart, *Organometallics* **1987**, *6*, 2484-2488; c) C. Scriban, D. S. Glueck, A. G. DiPasquale, A. L. Rheingold, *Organometallics* **2006**, *25*, 5435-5448; d) T. W. Chapp, A. J. Schoenfeld, M. D. Sanderson, C. L. Chang, D. S. Glueck, J. A. Golen, C. E. Moore, A. L. Rheingold, *Organometallics* **2012**, *31*, 5573-5585.
- [37] H. Schäfer, D. Binder, *Z. Anorg. Allg. Chem.* **1988**, *560*, 65-79.
- [38] The same species still formed even with up to 4 equiv PArH_2 and independent of the reaction temperature (-78 to 100 °C).
- [39] D. K. Wicht, S. N. Paisner, B. M. Lew, D. S. Glueck, G. P. A. Yap, L. M. Liable-Sands, A. L. Rheingold, C. M. Haar, S. P. Nolan, *Organometallics* **1998**, *17*, 652-660.
For NMR data on the free phosphines, see: J. W. Akitt, R. H. Cragg, N. N. Greenwood, *Chem. Commun.* **1966**, *5*, 134-135.
- [40] D. S. Bohle, G. R. Clark, C. E. F. Rickard, W. R. Roper, M. J. Taylor, *J. Organomet. Chem.* **1988**, *348*, 385-409.
- [41] The minor product was the dinuclear, P_2 -bridged complex $[\{\text{Ni}(\text{I}^i\text{Pr}_2)_2\}_2(\mu, \eta^{2:2}-\text{P}_2)]$ which was reported previously from the reaction of $[\text{Ni}_2(\text{I}^i\text{Pr}_2)_4(\text{COD})]$ with P_4 . B. Zarzycki, T. Zell, D. Schmidt, U. Radius. *Eur. J. Inorg. Chem.* **2013**, 2051-2058.
- [42] a) D. S. Bohle, T. C. Jones, C. E. F. Rickard, W. R. Roper, *Organometallics* **1986**, *5*, 1612-1619; b) A. J. Deeming, S. Doherty, J. E. Marshall, J. L. Powell, A. M. Senior, *J. Chem. Soc., Dalton Trans.* **1993**, 1093-1100; c) M. Driess, H. Pritzkow, U. Winkler, *J.*

- Organomet. Chem.* **1997**, 529, 313-321; d) A. K. Hickey, I. Muñoz, S. B., S. A. Lutz, M. Pink, C. H. Chen, J. M. Smith, *Chem. Commun.* **2017**, 53, 412-415; e) M. E. Garner, J. Arnold, *Organometallics* **2017**, 36, 4511-4514; f) D. Wang, Q. Chen, X. Leng, L. Deng, *Inorg. Chem.* **2018**, 57, 15600-15609; g) T. M. Rookes, E. P. Wildman, G. Balazs, B. M. Gardner, A. J. Wooles, M. Gregson, F. Tuna, M. Scheer, S. T. Liddle, *Angew. Chem. Int. Ed.* **2018**, 57, 1332-1336; *Angew. Chem.* **2018**, 130, 1346-1350.
- [43] D. S. Bohle, G. R. Clark, C. E. F. Rickard, W. R. Roper, *J. Organomet. Chem.* **1990**, 393, 243-285.
- [44] S. Gómez-Ruiz, E. Hey-Hawkins, *Dalton Trans.* **2007**, 5678-5683.
- [45] J. W. Ho, T. L. Breen, A. Ozarowski, D. W. Stephan, *Inorg. Chem.* **1994**, 33, 865-870.
- [46] J. Chatt, P. B. Hitchcock, A. Pidcock, C. P. Warrens, K. R. Dixon, *J. Chem. Soc., Dalton Trans.* **1984**, 2237-2244.
- [47] For computational studies that consider phosphine dissociation, see: a) Y. Zhao, D. G. Truhlar, *Org. Lett.* **2007**, 9, 1967-1970; b) Y. Minenkov, G. Occhipinti, V. R. Jensen, *J. Phys. Chem. A* **2009**, 113, 11833-11844; c) D. Benitez, E. Tkatchouk, W. A. Goddard, *Organometallics* **2009**, 28, 2643-2645; d) N. Sieffert, M. Bühl, *Inorg. Chem.* **2009**, 48, 4622-4624; e) C. L. McMullin, J. Jover, J. N. Harvey, N. Fey, *Dalton Trans.* **2010**, 39, 10833-10836; f) M. S. G. Ahlquist, P.-O. Norrby, *Angew. Chem. Int. Ed.* **2011**, 50, 11794-11797; *Angew. Chem.* **2011**, 123, 11998-12001.
- [48] For recent related examples at nickel see: a) L. V. Hooker, S. R. Neufeldt, *Tetrahedron* **2018**, 74, 6717-6725; b) E. D. Entz, J. E. A. Russell, L. V. Hooker, S. R. Neufeldt, *J. Am. Chem. Soc.* **2020**, 142, 15454-15463.
- [49] For recent reviews on the challenge of accuracy in computational chemistry, see: a) L. Perrin, K. J. T. Carr, D. McKay, C. L. McMullin, S. A. Macgregor, O. Eisenstein,

- Struct. Bond.* **2016**, *167*, 1-37; b) S. Grimme, P. R. Schreiner, *Angew. Chem. Int. Ed.* **2018**, *57*, 4170-4176; *Angew. Chem.* **2004**, *130*, 4241-4248; c) J. N. Harvey, F. Himo, F. Maseras, L. Perrin, *ACS Catal.* **2019**, *9*, 6803-6813.
- [50] a) A second PPh₂H ligand can add to **I** to give [Ni(Ime₄)₂(PPh₂H)₂], **IVb**, at -5.5 kcal/mol. As this lies below the energy of **5**, the observed product of P-H activation, the relative stability of **IVb** appears to be over-estimated with B97D and for the purposes of the discussion we have disregarded this. This energy difference is very functional dependent, but those functionals that place **IVb** higher in energy than **5** also gave too high a barrier for the P-H activation reactions. Identifying one functional that can capture all the features of the experimental system is highly problematic. b) For a rare example of a [Ni(NHC)₂(PR₃)₂] species, see: A. L. Ostericher, K. M. Waldie, C. P. Kubiak, *ACS Catal.* **2018**, *8*, 9596-9603.
- [51] The energy of this triplet species is again functional dependent but the behaviour of the B97D functional is generally similar to that of TPSS that was shown in a recent study to produce reliable spin-state energetics in a range of Co(III) complexes. S. E. Neale, D. A. Pantazis, S. A. Macgregor, *Dalton Trans.* **2020**, *49*, 6478-6487. See Supporting Information for details.
- [52] The direct reaction of *trans*-**III** with PPh₂H is possible but was found to be less accessible, proceeding through a transition state at +27.9 kcal/mol (see Figure S79 in the Supporting Information).
- [53] A trigonal bipyramidal η²-H₂ complex, **Int(IV-V)** and a corresponding H₂ loss transition state, **TS(IV-V)2** were located on the SCF surface, but **TS(IV-V)2** falls below **Int(IV-V)** upon applying the energy corrections. See Figure S71 in the Supporting Information.

- [54] The free energy of dissociation of IMe_4 from *cis*-**V** is 20.5 kcal/mol, ruling out a dissociative isomerisation pathway.
- [55] Possible proton, H atom or hydride transfers from PPh_2H to **5** were also assessed for the second P-H activation, but these proved significantly higher in energy. See Figure S80 in the Supporting Information.
- [56] I. V. Kourkine, M. D. Sargent, D. S. Glueck, *Organometallics* **1998**, *17*, 125-127.
- [57] P-C bond activation in $[\text{Ni}(\text{IMe}_4)_2(\text{PPh}_3)]$ to form **1** was also computed and found to proceed with an overall barrier of 23.1 kcal/mol. See Figure S77 in the Supporting Information.
- [58] The triplet form of **VI** lies at +18.5 kcal/mol and a transition state for P-H activation on the triplet surface was located at +21.3 kcal/mol. This leads directly to ^3V , but this process is significantly higher than the reaction on the singlet surface.
- [59] For experimental determinations, see: a) *Ru*: L. Jafarpour, S. P. Nolan, *J. Organomet. Chem.* **2001**, *617-618*, 17-27; b) *Ru*: A. C. Hillier, W. J. Sommer, B. S. Yong, J. L. Petersen, L. Cavallo, S. P. Nolan, *Organometallics* **2003**, *22*, 4322-4326; c) *Ni*: R. Dorta, E. D. Stevens, N. M. Scott, C. Costabile, L. Cavallo, C. D. Hoff, S. P. Nolan, *J. Am. Chem. Soc.* **2005**, *127*, 2485-2495.

[Ni(NHC)₂] as a Scaffold for Structurally Characterized *trans* [H-Ni-PR₂] and *trans* [R₂P-Ni-PR₂] Complexes

Sara Sabater, David Schmidt, Heidi Schmidt, nee Schneider, Maximilian W. Kuntze-Fechner, Thomas Zell, Connie J. Isaac, Nasir A. Rajabi, Harry Grieve, William J. M. Blackaby, John P. Lowe, Stuart A. Macgregor, Mary F. Mahon, Udo Radius and Michael K. Whittlesey

Whole Lotta Terminal Phosphidos: A combined experimental and computational study on the reactivity of [Ni(NHC)₂] (NHC = N-heterocyclic carbene) towards secondary and primary phosphines has been undertaken. P-H bond activation, as well as P-P activation of diphosphines, yields an extensive series of structurally characterised terminal phosphido complexes which exhibit unusual *trans* phosphido hydrido and novel *trans* bis-phosphido geometries.

